

Inverse scattering problem for a hyperbolic system of first order equations on a semi-axis on a first approximation

Lala N. Atamova · Aynur N. Safarova

Received: . 17.04.2019 / Revised: 23.07.2019 / Accepted: 04.11.2019

Abstract. In this paper for a hyperbolic system of five equations on the semi-axis, by joint consideration of three problems an inverse scattering problem on a first approximation was solved. The coefficients of the considered system are uniquely determined by the scattering operator on the semi-axis.

Keywords. inverse problem, scattering operator, factorization.

Mathematics Subject Classification (2010): 35J25; 35R30, 53C15

1 Introduction

Inverse scattering problems for different linear systems of first order hyperbolic equations on the axis and semi-axis were studied in the papers of L.P. Nizhnik [3], L.P. Nizhnik and V.G. Tarasov [2], A.S. Fokas and L.Y. Sung [1], N.Sh. Iskenderov [4], M.I. Ismailov [5] and others.

In this paper we study direct and inverse scattering problems for a system of five hyperbolic equations of first order on a semi-axis in the case when there are two given incident waves.

When there are three incident and two scattering waves, these problems were studied in [8] when there are four incident and two scattering waves, in [7].

2 Scattering problem on a semi-axis

On the semi-axis $x \geq 0$ consider a system of equations of the form:

$$\xi_i \frac{\partial U_i(x, t)}{\partial t} - \frac{\partial U_i(x, t)}{\partial x} = \sum_{j=1}^5 c_{ij}(x, t) U_j(x, t), i = \overline{1, 5} \quad (2.1)$$

L.N. Atamova
Khazar University, Baku, Azerbaijan
E-mail: LJafarova@khazar.org

A.N. Safarova
Institute of Mathematics and Mechanics of NAS of Azerbaijan, Baku, Azerbaijan
E-mail: safarova-aynur@bk.ru

where $c_{ij}(x, t)$ are complex-valued measurable functions with respect to x and t satisfying the conditions:

$$|c_{ij}(x, t)| \leq C [(1 + |x|)(1 + |t|)]^{-1-\varepsilon}, \quad (2.2)$$

moreover

$$c_{ii}(x, t) = 0, i = \overline{1, 5}, \xi_1 > \xi_2 > 0 > \xi_3 > \xi_4 > \xi_5, -\infty < t < +\infty$$

Let us consider system (2.1) on a semi-axis under three different boundary conditions:

$$1) \quad \begin{cases} U_3^1(0, t) = U_1^1(0, t) + U_2^1(0, t) \\ U_4^1(0, t) = U_2^1(0, t) \\ U_5^1(0, t) = U_1^1(0, t) \end{cases} \quad (2.3)$$

$$2) \quad \begin{cases} U_3^2(0, t) = U_1^2(0, t) \\ U_4^2(0, t) = U_1^2(0, t) + U_2^2(0, t) \\ U_5^2(0, t) = U_2^2(0, t) \end{cases} \quad (2.4)$$

$$3) \quad \begin{cases} U_3^3(0, t) = U_2^3(0, t) \\ U_4^3(0, t) = U_1^3(0, t) \\ U_5^3(0, t) = U_1^3(0, t) + U_2^3(0, t) \end{cases} \quad (2.5)$$

Any essentially bounded solution $U(x, t) = \{U_1(x, t), U_2(x, t), \dots, U_5(x, t)\}$ of the system (2.1) with the coefficients $c_{ij}(x, t), i, j = \overline{1, 5}$, satisfying conditions (2.2) admit on the semi-axis $x \geq 0$ the following asymptotic representations:

$$\begin{cases} U_i(x, t) = a_i(t + \xi_i x) + o(1), i = 1, 2 \\ U_i(x, t) = b_i(t + \xi_i x) + o(1), i = \overline{3, 5}, x \rightarrow +\infty, \end{cases} \quad (2.6)$$

where $a_i(s) \in L_\infty(-\infty, +\infty)$ ($i = 1, 2$) determine the incident waves, while $b_i(s) \in L_\infty(-\infty, +\infty), i = \overline{3, 5}$ the scattering ones.

The scattering problem for system (2.1) is in finding the solution to the system (2.1) by the given incident waves and boundary conditions for $x = 0$.

The scattering problem under joint consideration of the first, second and third problems is stated as follows: by the given function $a_1(s), a_2(s) \in L_\infty(R), R = (-\infty, +\infty)$ find the solution

$$U^k(x, t) \in L_\infty((0, +\infty) \times (-\infty, +\infty), C^2), (k = 1, 2)$$

of the first, second and third problems for which in L_∞ the following asymptotic representations are valid:

$$U_i^k(x, t) = a_i(t + \xi_i x) + o(1), x \rightarrow \infty, i = 1, 2, k = \overline{1, 3};$$

where $U^k(x, t) = (U_1^k(x, t), \dots, U_5^k(x, t))$.

Theorem 2.1 *Let the coefficients $c_{ij}(x, t), i, j = 5$ of the system (2.1), satisfy conditions (2.2). Then there exists a unique solution of the scattering problem on the semi-axis $x \geq 0$ for the system (2.1) with arbitrary given incident waves*

$$a_i(s) \in L_\infty(R), R = (-\infty, +\infty), i = 1, 2$$

The proof of this theorem is similar to one in [7].

Note that the scattering problem for the k -th ($k = \overline{1, 3}$) problem is equivalent to the following system of integral equations:

$$\begin{aligned} U_1^k(x, t) &= a_1(t + \xi_1 x) + \int_x^{+\infty} \sum_{j=1}^5 (c_{1j} U_j)(y, t + \xi_1(x - y)) dy, \\ U_2^k(x, t) &= a_2(t + \xi_2 x) + \int_x^{+\infty} \sum_{j=1}^5 (c_{2j} U_j)(y, t + \xi_2(x - y)) dy, \\ U_i^k(x, t) &= b_i(t + \xi_i x) + \int_x^{+\infty} \sum_{j=1}^5 (c_{ij} U_j)(y, t + \xi_i(x - y)) dy, i = \overline{3, 5}, \end{aligned} \quad (2.7)$$

where the functions $b_3^k(s), b_4^k(s), b_5^k(s)$, $k = 1, 2$ are expressed by $a_1(s), a_2(s)$ the coefficients $c_{ij}(x, t)$, $i, j = \overline{1, 5}$ and the solutions of the first, second and third problems, respectively, in the following way:

$$\left\{ \begin{array}{l} b_3^1(t) = a_1(t) + a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^1(y, t - \xi_1 y) + \\ + c_{2j}(y, t - \xi_2 y) U_j^1(y, t - \xi_2 y) - c_{3j}(y, t - \xi_3 y) U_j^1(y, t - \xi_3 y)] dy, \\ b_4^1(t) = a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{2j}(y, t - \xi_2 y) U_j^1(y, t - \xi_2 y) - c_{4j}(y, t - \xi_4 y) U_j^1(y, t - \xi_4 y)] dy, \\ b_5^1(t) = a_1(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^1(y, t - \xi_1 y) - \\ - c_{5j}(y, t - \xi_5 y) U_j^1(y, t - \xi_5 y)] dy \end{array} \right. \quad (2.8)$$

$$\left\{ \begin{array}{l} b_3^2(t) = a_1(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^2(y, t - \xi_1 y) - \\ - c_{3j}(y, t - \xi_3 y) U_j^2(y, t - \xi_3 y)] dy, \\ b_4^2(t) = a_1(t) + a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^2(y, t - \xi_1 y) + \\ + c_{2j}(y, t - \xi_2 y) U_j^2(y, t - \xi_2 y) - c_{4j}(y, t - \xi_4 y) U_j^2(y, t - \xi_4 y)] dy, \\ b_5^2(t) = a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{2j}(y, t - \xi_2 y) U_j^2(y, t - \xi_2 y) - \\ - c_{5j}(y, t - \xi_5 y) U_j^2(y, t - \xi_5 y)] dy \end{array} \right. \quad (2.9)$$

$$\left\{ \begin{array}{l} b_3^3(t) = a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{2j}(y, t - \xi_2 y) U_j^3(y, t - \xi_2 y) - \\ - c_{3j}(y, t - \xi_3 y) U_j^3(y, t - \xi_3 y)] dy, \\ b_4^3(t) = a_1(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^3(y, t - \xi_1 y) - \\ - c_{4j}(y, t - \xi_4 y) U_j^3(y, t - \xi_4 y)] dy \\ b_5^3(t) = a_1(t) + a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^3(y, t - \xi_1 y) + \\ + c_{2j}(y, t - \xi_2 y) U_j^3(y, t - \xi_2 y) - c_{5j}(y, t - \xi_5 y) U_j^3(y, t - \xi_5 y)] dy \end{array} \right. \quad (2.10)$$

It follows from theorem (2.1) that to each vector-function $a(t) = (a_1(t), a_2(t)) \in L_\infty(R)$ giving the incident waves there correspond the solutions of three scattering problems of the system (2.1) with boundary conditions (2.3), (2.4), (2.5) and the given asymptotics

$$\begin{cases} U_3^k(x, t) = b_3^k(t + \xi_3 x) + o(1), \\ U_4^k(x, t) = b_4^k(t + \xi_4 x) + o(1), \\ U_5^k(x, t) = b_5^k(t + \xi_5 x) + o(1), k = \overline{1, 3}, x \rightarrow +\infty, \end{cases} \quad (2.11)$$

i.e. the vector of scattering waves $b(t) = (b^1(t), b^2(t), b^3(t))$, where $b^k(t) = (b_3^k(t), b_4^k(t), b_5^k(t))$ ($k = \overline{1, 3}$). Relation (2.11) follows from (2.7) and conditions (2.2). Thus, in the space of essentially bounded functions, we determined the operator $S = (S^1, S^2, S^3)$ that takes $a(t)$ to $b(t)$:

$$S^k \begin{pmatrix} a_1(t) \\ a_2(t) \end{pmatrix} = \begin{pmatrix} b_3^k(t) \\ b_4^k(t) \\ b_5^k(t) \end{pmatrix}, k = \overline{1, 3} \quad (2.12)$$

Here,

$$S = (S^1, S^2, S^3) \text{ and } S^k = \begin{pmatrix} S_{11}^k S_{12}^k \\ S_{21}^k S_{22}^k \\ S_{31}^k S_{32}^k \end{pmatrix}, k = \overline{1, 3}$$

$$\begin{aligned} b_3^k(t) &= S_{11}^k a_1(t) + S_{12}^k a_2(t), \\ b_4^k(t) &= S_{21}^k a_1(t) + S_{22}^k a_2(t), \\ b_5^k(t) &= S_{31}^k a_1(t) + S_{32}^k a_2(t), k = \overline{1, 3}. \end{aligned} \quad (2.13)$$

From (2.8), (2.9), (2.10) it follows that the elements of the operator have S^k ($k = \overline{1, 3}$) have the form:

$$\begin{cases} S_{11}^1 = I + F_{11}^1, S_{12}^1 = I + F_{12}^1, \\ S_{21}^1 = F_{21}^1, S_{22}^1 = I + F_{22}^1, \\ S_{31}^1 = I + F_{31}^1, S_{32}^1 = F_{32}^1, \\ \\ S_{11}^2 = I + F_{11}^2, S_{12}^2 = F_{12}^2, \\ S_{21}^2 = I + F_{21}^2, S_{22}^2 = I + F_{22}^2, \\ S_{31}^2 = F_{32}^2, S_{32}^2 = I + F_{32}^2, \\ \\ S_{11}^3 = F_{11}^3, S_{12}^3 = I + F_{12}^3, \\ S_{21}^3 = I + F_{21}^3, S_{22}^3 = F_{22}^3, \\ S_{31}^3 = I + F_{31}^3, S_{32}^3 = I + F_{32}^3, \end{cases} \quad (2.14)$$

where the operators F_{ij}^k ($j = 1, 2, i, k = \overline{1, 3}$) are Fredholm integral operators.

3 The inverse scattering problem on a semi-axis on a first approximation

The inverse problem for the system (2.1), is in finding the coefficients of the system (2.1), by the given scattering operator S on a semi-axis.

Here the coefficients of the system (2.1), are restored by the scattering operator on a semi-axis constructed on a first approximation. It is constructed in the explicit form.

For the first problem as a zero order approximation we take

$$\begin{aligned} U_k^{(1,0)}(x, t) &= a_k(t + \xi_k x), k = 1, 2, \\ U_3^{(1,0)}(x, t) &= a_1(t + \xi_3 x) + a_2(t + \xi_3 x), \\ U_4^{(1,0)}(x, t) &= a_2(t + \xi_4 x), \\ U_5^{(1,0)}(x, t) &= a_1(t + \xi_5 x); \end{aligned}$$

for the second problem

$$\begin{aligned} U_k^{(2,0)}(x, t) &= a_k(t + \xi_k x), k = 1, 2, \\ U_3^{(2,0)}(x, t) &= a_1(t + \xi_3 x), \\ U_4^{(2,0)}(x, t) &= a_1(t + \xi_4 x) + a_2(t + \xi_4 x), \\ U_5^{(2,0)}(x, t) &= a_2(t + \xi_5 x); \end{aligned}$$

for the third problem

$$\begin{aligned} U_k^{(3,0)}(x, t) &= a_k(t + \xi_k x), k = 1, 2, \\ U_3^{(3,0)}(x, t) &= a_2(t + \xi_3 x), \\ U_4^{(3,0)}(x, t) &= a_1(t + \xi_4 x), \\ U_5^{(3,0)}(x, t) &= a_1(t + \xi_5 x) + a_2(t + \xi_5 x); \end{aligned}$$

Then in equalities (2.8), (2.9)and (2.10) the first order approximations will be:

$$\left\{ \begin{array}{l} b_3^1(t) = a_1(t) + a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y)U_j^{(1,0)}(y, t - \xi_1 y) + \\ + c_{2j}(y, t - \xi_2 y)U_j^{(1,0)}(y, t - \xi_2 y) - c_{3j}(y, t - \xi_3 y)U_j^{(1,0)}(y, t - \xi_3 y)]dy, \\ b_4^1(t) = a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{2j}(y, t - \xi_2 y)U_j^{(1,0)}(y, t - \xi_2 y) - \\ - c_{4j}(y, t - \xi_4 y)U_j^{(1,0)}(y, t - \xi_4 y)]dy, \\ b_5^1(t) = a_1(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y)U_j^{(1,0)}(y, t - \xi_1 y) - \\ - c_{5j}(y, t - \xi_5 y)U_j^{(1,0)}(y, t - \xi_5 y)]dy; \end{array} \right. \quad (3.1)$$

$$\left\{ \begin{array}{l} b_3^2(t) = a_1(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y)U_j^{(2,0)}(y, t - \xi_1 y) - \\ - c_{3j}(y, t - \xi_3 y)U_j^{(2,0)}(y, t - \xi_3 y)]dy, \\ b_4^2(t) = a_1(t) + a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y)U_j^{(2,0)}(y, t - \xi_1 y) + \\ + c_{2j}(y, t - \xi_2 y)U_j^{(2,0)}(y, t - \xi_2 y) - \\ - c_{4j}(y, t - \xi_4 y)U_j^{(2,0)}(y, t - \xi_4 y)]dy, \\ b_5^2(t) = a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{2j}(y, t - \xi_2 y)U_j^{(2,0)}(y, t - \xi_2 y) - \\ - c_{5j}(y, t - \xi_5 y)U_j^{(2,0)}(y, t - \xi_5 y)]dy; \end{array} \right. \quad (3.2)$$

$$\left\{ \begin{array}{l} b_3^3(t) = a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{2j}(y, t - \xi_2 y) U_j^{(3,0)}(y, t - \xi_2 y) - \\ - c_{3j}(y, t - \xi_3 y) U_j^{(3,0)}(y, t - \xi_3 y)] dy, \\ b_4^3(t) = a_1(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^{(3,0)}(y, t - \xi_1 y) - \\ - c_{4j}(y, t - \xi_4 y) U_j^{(3,0)}(y, t - \xi_4 y)] dy, \\ b_5^3(t) = a_1(t) + a_2(t) + \int_0^{+\infty} \sum_{j=1}^5 [c_{1j}(y, t - \xi_1 y) U_j^{(3,0)}(y, t - \xi_1 y) + \\ + c_{2j}(y, t - \xi_2 y) U_j^{(3,0)}(y, t - \xi_2 y) - c_{5j}(y, t - \xi_5 y) U_j^{(3,0)}(y, t - \xi_5 y)] dy; \end{array} \right. \quad (3.3)$$

respectively.

Taking into account in (2.13) and (2.14), we have:

$$\begin{aligned} c_{12}(x, y) &= (\xi_2 - \xi_1) [F_{12}^1(\xi_1 x + y, \xi_2 x + y) - F_{12}^2(\xi_1 x + y, \xi_2 x + y)], \\ c_{13}(x, y) &= (\xi_1 - \xi_3) [F_{12}^2(\xi_1 x + y, \xi_3 x + y) - F_{12}^1(\xi_1 x + y, \xi_3 x + y) + \\ &+ \frac{1}{2} [F_{22}^3(\xi_1 x + y, \xi_3 x + y) + F_{32}^1(\xi_1 x + y, \xi_3 x + y) - F_{21}^3(\xi_1 x + y, \xi_3 x + y)]] \\ c_{14}(x, y) &= \frac{\xi_1 - \xi_4}{2} [F_{21}^3(\xi_1 x + y, \xi_4 x + y) - F_{22}^3(\xi_1 x + y, \xi_4 x + y) \\ &\quad + F_{32}^1(\xi_1 x + y, \xi_4 x + y)], \\ c_{15}(x, y) &= (\xi_1 - \xi_5) [F_{31}^1(\xi_1 x + y, \xi_5 x + y) - F_{12}^2(\xi_3 x + y, \xi_5 x + y) + F_{12}^1(\xi_3 x + y, \xi_5 x + y) + \\ &+ \frac{1}{2} [F_{21}^3(\xi_1 x + y, \xi_5 x + y) - F_{22}^3(\xi_1 x + y, \xi_5 x + y) - F_{32}^1(\xi_1 x + y, \xi_5 x + y)]], \\ c_{21}(x, y) &= (\xi_1 - \xi_2) [F_{11}^1(\xi_2 x + y, \xi_1 x + y) - F_{11}^2(\xi_2 x + y, \xi_1 x + y)], \\ c_{23}(x, y) &= (\xi_2 - \xi_3) [F_{22}^1(\xi_2 x + y, \xi_3 x + y) - F_{11}^2(\xi_2 x + y, \xi_3 x + y) + \\ &\quad + F_{12}^1(\xi_2 x + y, \xi_3 x + y)], \\ c_{24}(x, y) &= (\xi_4 - \xi_2) [F_{11}^2(\xi_2 x + y, \xi_4 x + y) - F_{12}^1(\xi_2 x + y, \xi_4 x + y)], \\ c_{25}(x, y) &= (\xi_2 - \xi_5) [F_{32}^2(\xi_2 x + y, \xi_5 x + y) - F_{11}^2(\xi_2 x + y, \xi_5 x + y) + \\ &\quad + F_{12}^1(\xi_2 x + y, \xi_5 x + y)], \\ c_{31}(x, y) &= (\xi_3 - \xi_1) F_{11}^2(\xi_3 x + y, \xi_1 x + y), \\ c_{32}(x, y) &= (\xi_3 - \xi_2) F_{12}^2(\xi_3 x + y, \xi_2 x + y), \\ c_{34}(x, y) &= (\xi_4 - \xi_3) [F_{11}^3(\xi_3 x + y, \xi_4 x + y) - F_{12}^3(\xi_3 x + y, \xi_4 x + y) + 2F_{12}^1(\xi_3 x + y, \xi_4 x + y) - \\ &\quad - 2F_{11}^2(\xi_3 x + y, \xi_4 x + y) - F_{22}^1(\xi_3 x + y, \xi_4 x + y)], \\ c_{35}(x, y) &= (\xi_5 - \xi_3) [F_{12}^3(\xi_3 x + y, \xi_5 x + y) - 2F_{12}^1(\xi_3 x + y, \xi_5 x + y) + 2F_{11}^2(\xi_3 x + y, \xi_5 x + y) - \\ &\quad - F_{32}^2(\xi_3 x + y, \xi_5 x + y) - F_{22}^1(\xi_3 x + y, \xi_5 x + y)], \\ c_{41}(x, y) &= (\xi_4 - \xi_1) F_{21}^3(\xi_4 x + y, \xi_1 x + y), \end{aligned}$$

$$\begin{aligned}
c_{42}(x, y) &= (\xi_4 - \xi_2) F_{22}^2(\xi_4 x + y, \xi_2 x + y), \\
c_{43}(x, y) &= (\xi_4 - \xi_3) [F_{22}^1(\xi_4 x + y, \xi_3 x + y) - F_{22}^2(\xi_4 x + y, \xi_3 x + y)], \\
c_{45}(x, y) &= (\xi_5 - \xi_4) [F_{22}^2(\xi_4 x + y, \xi_5 x + y) - F_{31}^3(\xi_4 x + y, \xi_5 x + y) - F_{12}^1(\xi_4 x + y, \xi_5 x + y) + \\
&\quad + F_{12}^2(\xi_4 x + y, \xi_5 x + y)], \\
c_{51}(x, y) &= (\xi_5 - \xi_1) [F_{31}^1(\xi_5 x + y, \xi_1 x + y) + F_{32}^2(\xi_5 x + y, \xi_1 x + y) - F_{32}^1(\xi_5 x + y, \xi_1 x + y)], \\
c_{52}(x, y) &= (\xi_5 - \xi_2) [F_{32}^3(\xi_5 x + y, \xi_2 x + y) - F_{32}^1(\xi_5 x + y, \xi_2 x + y) + F_{32}^2(\xi_5 x + y, \xi_2 x + y)], \\
c_{53}(x, y) &= (\xi_5 - \xi_3) [F_{32}^1(\xi_5 x + y, \xi_3 x + y) - F_{32}^2(\xi_5 x + y, \xi_3 x + y)], \\
c_{54}(x, y) &= (\xi_5 - \xi_4) [F_{32}^1(\xi_5 x + y, \xi_4 x + y) - F_{32}^3(\xi_5 x + y, \xi_4 x + y)]. \tag{3.4}
\end{aligned}$$

where $F_{ij}^k(t, \tau)$ -kernel of the operators $F_{ij}^k(j = 1, 2, k = \overline{1, 3})$. Note that from the remaining relations we have:

$$\begin{aligned}
F_{21}^1(t, \tau) &= F_{21}^2(t, \tau), \tau > t, \\
F_{11}^1(t, \tau) &= F_{31}^1(t, \tau) + F_{12}^3(t, \tau), \tau < t, \\
F_{11}^1(t, \tau) &= F_{12}^3(t, \tau) + F_{12}^2(t, \tau) - F_{12}^1(t, \tau) + F_{31}^1(t, \tau), \tau < t, \\
F_{12}^1(t, \tau) &= F_{11}^3(t, \tau) - F_{12}^3(t, \tau) - F_{11}^2(t, \tau) - F_{22}^1(t, \tau) + \\
&\quad + F_{12}^2(t, \tau) + F_{32}^1(t, \tau), \tau < t, \\
F_{21}^1(t, \tau) &= F_{22}^3(t, \tau) - F_{31}^3(t, \tau) + F_{12}^2(t, \tau) + F_{22}^1(t, \tau) - \\
&\quad - 2F_{11}^2(t, \tau) + F_{32}^2(t, \tau) + F_{12}^1(t, \tau), \tau < t, \\
F_{11}^2(t, \tau) &= F_{11}^3(t, \tau) - F_{12}^3(t, \tau) + F_{12}^1(t, \tau) - 2F_{11}^2(t, \tau) - \\
&\quad - F_{22}^1(t, \tau) + F_{12}^2(t, \tau) + F_{32}^1(t, \tau), \tau < t, \\
F_{12}^2(t, \tau) &= F_{11}^3(t, \tau) - 2F_{22}^1(t, \tau) - F_{32}^2(t, \tau) + 2F_{12}^1(t, \tau) - 2F_{12}^2(t, \tau) + \\
&\quad + F_{21}^3(t, \tau) - F_{22}^3(t, \tau) + F_{31}^1(t, \tau), \tau < t, \\
F_{21}^2(t, \tau) &= F_{12}^2(t, \tau) + F_{32}^1(t, \tau) - F_{12}^1(t, \tau) + F_{22}^1(t, \tau), \tau < t, \\
F_{21}^2(t, \tau) &= F_{21}^3(t, \tau) + F_{22}^1(t, \tau) - F_{22}^2(t, \tau) + F_{11}^1(t, \tau) - F_{11}^2(t, \tau), \tau > t \\
F_{31}^3(t, \tau) &= F_{21}^3(t, \tau) - F_{22}^3(t, \tau) + F_{31}^1(t, \tau) - F_{12}^2(t, \tau) - \\
&\quad - F_{12}^1(t, \tau) + F_{32}^2(t, \tau), \tau < t, \tag{3.5} \\
F_{31}^2(t, \tau) &= F_{31}^1(t, \tau) + F_{32}^1(t, \tau) - F_{32}^3(t, \tau) + F_{11}^1(t, \tau) - F_{11}^2(t, \tau), \tau > t, \\
F_{11}^3(t, \tau) &= F_{11}^1(t, \tau), \tau > t, \\
F_{12}^3(t, \tau) &= F_{12}^2(t, \tau), \tau < t \\
F_{22}^3(t, \tau) &= F_{22}^1(t, \tau), \tau > t, \\
F_{31}^3(t, \tau) &= F_{31}^1(t, \tau) + F_{32}^2(t, \tau) - F_{32}^3(t, \tau) + F_{11}^1(t, \tau) - F_{11}^2(t, \tau), \tau > t \\
F_{32}^3(t, \tau) &= F_{12}^1(t, \tau) + F_{31}^1(t, \tau) - F_{12}^2(t, \tau), \tau < t,
\end{aligned}$$

The scattering operator S has 18 elements on the axis or 36 elements on the semi-axis. From 36 elements by formula (3.4) we find 20 coefficients of the system (2.1), 16 unnecessary elements are connected with 8 relations on the axis (16 on the semi-axis with respect to t) by formula (3.5).

References

1. Fokas, A.S., Sung, L.Y.: On the solvability of the N-wave, Davey- Stewartson and Kadomstev -Petsiasvili equations. *Inverse Problems*, 8: 673-708 (1992).
2. Nizhnik L.P., Tarasov V.G.: Inverse non-stationary scattering problem for hyperbolic system of equations, *DAN SSSR*, **233** (3): 300-303 (1977) (in Russian).
3. Nizhnik, L.P.: Inverse scattering problems for hyperbolic equations, *Kiev. Nauk. Dumka*, 232 p. (1991).
4. Iskenderov, N.Sh.: *Inverse scattering problem for a hyperbolic system of n equations of first order on a semi-axis*, Ukr. Mat. Zh. 43(12): 1638-1646 (1991).
5. Ismailov, M.I.: *Inverse scattering problem for hyperbolic systems on a semi-axis in the case of equal number of incident and scattered waves*, Inverse Problem, **22**: 955-974 (2006).
6. Iskenderov, N.Sh., Jabbarova, K.A.: *The inverse scattering problem for the system of five first order hyperbolic equations on semi-axis*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Math and Mech. (7), 41-54 (2005).
7. Iskenderov, N.Sh., Jafarova L.N.: *Direct and inverse scattering problems for the system of six first order hyperbolic equations on semi-axis*, Trans. Inst. Math. NAS Ukr. **14** (3), 67-99 (2017).
8. Mamedov, A.A.: *Scattering problem on a semi-axis for a system of first order hyperbolic equations*, Bull. Baku State Univ. **1**, 59-65 (2015).
9. Nizhnik, L.P., Tarasov, V.G.: *Inverse scattering problem for one-velocity transfer equation*, Dokl. AN SSSR. **242** (6), 1307-1310 (1978).